

SCHRÖDER'S PROBLEMS AND SCALING LIMITS OF RANDOM TREES

JIM PITMAN AND DOUGLAS RIZZOLO

ABSTRACT. In his now classic paper [17], Schröder posed four combinatorial problems about the number of certain types of bracketings of words and sets. Here we address what these bracketings look like on average. For each of the four problems we prove that a uniform pick from the appropriate set of bracketings, when considered as a tree, has the Brownian continuum random tree as its scaling limit as the size of the word or set goes to infinity.

1. INTRODUCTION

In his now classic paper [17], Schröder posed four combinatorial problems about bracketings of words and sets: how many binary bracketings are there of a word of length n ? how many bracketings are there of a word of length n ? how many binary bracketings are there of a set of size n ? and how many bracketings are there of a set of size n ? These questions are well studied and [18] gives a good account of the solutions. In this paper we are concerned with a probabilistic variation on these questions: for each of the above questions, if you select a bracketing uniformly at random what does it look like? To answer these questions, we will use the well known correspondence of bracketings described above to various types of trees. We will then apply Aldous's theory of continuum trees, originally developed in the series [1, 2, 3] and subsequently studied by many authors, to study the scaling limits of these trees. Let us briefly describe the correspondence between bracketings and trees.

The first problem: The correspondence is best illustrated by example. For $n = 4$ the binary word bracketings are


$$(xx)(xx) \quad x(x(xx)) \quad ((xx)x)x \quad x((xx)x) \quad (x(xx))x.$$

A binary bracketing of a word with n letters corresponds to rooted ordered binary tree with n leaves in a natural way. This is most easily described if we put brackets around the entire word and each letter, which are left out of our example because they are visually cumbersome. The tree corresponding to a bracketing is constructed recursively. A single bracketed letter is a leaf. For a word with more than one letter, the bracketing of the whole word is the root. Attached as subtrees to the root are, in order of appearance, the trees corresponding to the maximal proper bracketed subwords. For $n = 4$, this is illustrated by Figure 1.

It is worth noting that these trees are in bijection with rooted ordered trees with n vertices, but this correspondence is not as natural as the one above.


JP supported in part by NSF Grant No. 0806118.

DR supported in part by NSF Grant No. 0806118 and in part by the National Science Foundation Graduate Research Fellowship under Grant No. DGE 1106400.

FIGURE 1. Binary word bracketings and rooted ordered binary trees for $n = 4$


The second problem: General word bracketings are defined similarly to binary word bracketings and correspond to rooted ordered trees with n leaves and no vertices with out degree equal to one.

The third problem: The trees associated to binary set bracketings are constructed similarly to those associated to binary word bracketings. They are rooted, unordered, leaf-labeled binary trees. Figure 2 shows a sample of the correspondence for $n = 4$ (for $n = 4$ there are 15 bracketings, so showing the whole correspondence is unwieldy).

FIGURE 2. Binary set bracketings and rooted unordered leaf-labeled binary trees for $n = 4$

The fourth problem: General set bracketings are defined similarly to binary set bracketings and correspond to rooted unordered leaf-labeled trees with n leaves and no vertices with out degree equal to one. In the literature, these trees are also called fragmentation trees [9] and hierarchies [7]. The correspondence for $n = 3$ is in Figure 3.

Scaling limits of uniform picks from the trees appearing in the first and third problems are well studied. A uniform pick from rooted ordered binary tree's with n leaves has the same distribution as a Galton-Watson tree with offspring distribution $\xi_0 = \xi_2 = 1/2$ conditioned to have $2n - 1$ vertices. Thus it falls within the scope of the results in [3]. Similarly, a uniform pick from rooted unordered leaf-labeled binary tree's with n leaves is a uniform binary fragmentation tree with n leaves, and scaling limits of these are studied in [9]. In


 FIGURE 3. Set bracketings and rooted unordered leaf-labeled trees for $n = 3$

this paper we present a unified approach that is able to handle all four of these types of trees simultaneously. Our method is essentially to link the trees appearing in these four problems to Galton-Watson trees conditioned on their number of leaves. Scaling limits for these Galton-Watson trees were recently proven in [16]. We also draw a connection to certain Gibbs fragmentation trees, which were originally studied in [12]. Our main result is the following theorem, the notation for which will be fully explained later.

Theorem 1. *For $i = 1, 2, 3, 4$, let T_n^i be a uniform random tree of the type appearing in Schröder's i 'th problem with n leaves. For each i and n equip T_n^i with the graph metric where edges have length one and the uniform probability measure on its leaves. We then have the following limits with respect to the rooted Gromov-Hausdorff-Prokhorov topology:*

$$\begin{aligned}
 (i) \quad \frac{1}{\sqrt{n}} T_n^1 &\xrightarrow{d} 2\sqrt{2} T^{Br} & (ii) \quad \frac{1}{\sqrt{n}} T_n^2 &\xrightarrow{d} \frac{\sqrt{2}}{2\sqrt{\sqrt{2}-1}} T^{Br} \\
 (iii) \quad \frac{1}{\sqrt{n}} T_n^3 &\xrightarrow{d} 2\sqrt{2} T^{Br} & (iv) \quad \frac{1}{\sqrt{n}} T_n^4 &\xrightarrow{d} \frac{2}{\sqrt{4\log(2)-2}} T^{Br},
 \end{aligned}$$

where T^{Br} is the Brownian continuum random tree.

As noted above, parts (i) and (iii) were originally proven in [3] and [9] respectively. Parts (ii) and (iv) appear to be new.

The paper is organized as follows. In Section 2 we rigorously introduce the models of random trees under consideration here. In Section 3 we introduce the analytic setting for Theorem 1 and end with the proof of this theorem. Finally, in Section 4 we use elementary methods from analytic combinatorics to compute some asymptotic properties of these trees explicitly.

2. COMBINATORIAL MODELS AND GALTON-WATSON TREES

In this section we develop several combinatorial and probabilistic models of trees. There are two primary types of trees we will be dealing with in the sequel: rooted ordered unlabeled trees and rooted unordered leaf-labeled trees. Combinatorial relations between rooted ordered unlabeled trees and rooted unordered labeled trees are well known when the size of a tree is its number of vertices (see e.g. [15, 2, 7, 5]). In this section we develop analogous relations when the size of a tree is its number of leaves. Particularly important for us is Corollary 2, which relates Schröder's problems to particular Galton-Watson trees conditioned on their number of leaves.

We briefly give an account of the formal constructions of the trees we will be considering. Fix a countably infinite set S ; we will consider the vertex sets of all graphs discussed to be subsets of S . Let $\mathcal{T}_n$ denote the set of rooted unordered trees with n leaves (where the root is considered a leaf if and only if it is the only vertex in the tree) whose leaves are labeled by $\{1, 2, \dots, n\}$. More precisely, we consider the set $\mathcal{T}_n^S$ of all trees whose vertex sets are contained in S that have a distinguished root and n leaves (where the root is considered a leaf if and only if it is the only vertex in the tree), whose leaves are labeled by $\{1, 2, \dots, n\}$ and set $\mathcal{T}_n = \mathcal{T}_n^S / \sim$ where $t \sim s$ if there is a root and label preserving isomorphism from t to s . This is the only time we shall go through this formal construction, but all other sets of trees we discuss should be considered as formally constructed in an analogous fashion. We also let $\mathcal{T} = \cup_{n \geq 1} \mathcal{T}_n$. We let $\mathcal{T}_n^{(o)}$ be the set of rooted ordered unlabeled trees with n leaves and $\mathcal{T}^{(o)} = \cup_{n \geq 1} \mathcal{T}_n^{(o)}$.

We will be proving analogous results for trees in $\mathcal{T}$ and $\mathcal{T}^{(o)}$, so analogous that the only difference in the statements will be the superscript (o) . To avoid repetition we will use $\mathcal{T}^*$ and $\mathcal{T}_n^*$ when we don't want to specify whether we are in $\mathcal{T}$, $\mathcal{T}^{(o)}$, $\mathcal{T}_n$ or $\mathcal{T}_n^{(o)}$. That is, in a given result you may replace all the $*$'s by nothing or (o) . For a tree $t \in \mathcal{T}^*$, we define $|t|$ to be the number of leaves in t and $\#t$ to be the number of vertices in t .

2.1. Probabilities on trees. Let $\zeta = (\zeta_i)_{i \geq 0}$ be a sequence of numbers. We may then define the weight of a tree $t \in \mathcal{T}^*$ to be

$$w_\zeta(t) = \prod_{v \in t} \zeta_{\deg(v)}.$$

Here and throughout, $\deg(v)$ is the out degree of v , i.e., the number of children of v . We will assume the following conditions:

Condition 1. (i) $\zeta_i \geq 0$ for all i , (ii) $\zeta_0 > 0$, and (iii) for each n we have $\sum_{t \in \mathcal{T}_n^*} w_\zeta(t) < \infty$.

Observe that this condition is satisfied whenever $\zeta_1 = 0$, as is the case for Schröder's problems. For each n such that $w_\zeta(t) > 0$ for some $t \in \mathcal{T}_n^*$ we may define a probability measure on $\mathcal{T}_n^*$ by

$$Q_n^{\zeta^*}(t) = \frac{w_\zeta(t)}{\sum_{s \in \mathcal{T}_n^*} w_\zeta(s)}.$$

We wish to consider generating functions, but we want an ordinary generating function for $\mathcal{T}^{(o)}$ and an exponential generating function for $\mathcal{T}$. In order to do this all at once, for $z \in \mathbb{C}$, we define $y_n(z) = z^n/n!$ and $y_n^{(o)}(z) = z^n$, both for $n \geq 0$, and we use y_n^* in the same fashion as $\mathcal{T}^*$. The weighted generating function induced on $\mathcal{T}^*$ by ζ with the weights defined above is

$$C_\zeta^*(z) = \sum_{t \in \mathcal{T}^*} w_\zeta(t) y_{|t|}^*(z).$$

Letting $G_{\zeta,*}(z) = \sum_{i=1}^{\infty} \zeta_i y_i^*(z)$, is then easy to see that C_ζ^* satisfies the functional equation

$$(2.1) \quad C_\zeta^*(z) = \zeta_0 z + G_{\zeta,*}(C_\zeta^*(z)),$$

in the sense of formal power series. Our interest is in the measures Q_n^ζ and, in particular, we would like to find a Galton-Watson tree T such that Q_n^ζ is the law of T conditioned to

have n leaves. Recall that if $(\xi_i)_{i \geq 0}$ is a distribution on $\mathbb{Z}_+$ with mean less than or equal to one and $\xi_0 > 0$, a Galton-Watson tree with offspring distribution ξ is a random element T of $\mathcal{T}^{(o)}$ with law

$$\mathbb{P}(T = t) = \prod_{v \in t} \xi_{\deg(v)}.$$

T is called critical if ξ has mean equal to one. This leads to the notion of tilting, which is similar to exponential tilting for Galton-Watson trees conditioned on their number of vertices.

Proposition 1. *Suppose that ζ satisfies Condition 1 and suppose that $a, b > 0$. Define $\tilde{\zeta}$ by*

$$\tilde{\zeta}_0 = a\zeta_0 \quad \text{and} \quad \tilde{\zeta}_i = b^{i-1}\zeta_i \quad \text{for } i \geq 1.$$

Then $Q_n^{\zeta^*} = Q_n^{\tilde{\zeta}^*}$ for all $n \geq 1$.

Proof. This follows immediately from the computation that, for $t \in \mathcal{T}_n^*$, $w_{\tilde{\zeta}}(t) = a^n b^{n-1} w_{\zeta}(t)$. $\square$

The above result is the equivalent of exponential tilting for trees conditioned on their number of vertices. A consequence of this is that we can find a Galton-Watson tree T such that $Q_n^{\zeta^{(o)}}$ is the law of T conditioned to have n leaves if we can find $a, b > 0$ such that

$$a\zeta_0 + \frac{G_{\zeta, (o)}(b)}{b} = 1.$$

Furthermore, T will be critical if $G_{\zeta, (o)}'(b) = 1$. An immediate consequence of this is the following corollary.

Corollary 1. *Let $\xi^u = (\xi_i^u)_{i=0}^\infty$ be the probability distribution defined by*

$$\xi_0^u = \frac{2(3 - 2\sqrt{2})}{2 - \sqrt{2}} \approx 0.5858, \quad \xi_1^u = 0, \quad \text{and} \quad \xi_i^u = \left(\frac{2 - \sqrt{2}}{2} \right)^{i-1} \approx (0.2929)^{i-1} \quad \text{for } i \geq 2.$$

Note that ξ^u has mean 1 and variance $4\sqrt{2}$. Let T be a Galton-Watson tree with offspring distribution ξ^u . Then the law of T conditioned to have n leaves is uniform on the subset of $\mathcal{T}_n^{(o)}$ of trees with no vertices of out degree one.

Proof. The proof follows immediately from the discussion above by noting that, if $\zeta_i = 1$ for $i \neq 1$ and $\zeta_1 = 0$ then $Q_n^{\zeta^{(o)}}$ is uniform on $\mathcal{T}_n^{(o)}$. Explicitly, the distribution ξ^u is found by solving $G'_{\zeta, (o)}(b) = 1$, setting $a = (b - G_{\zeta, (o)}(b))/b$, and tilting as in Proposition 1. $\square$

Given the similarities in the construction of Q_n^ζ and $Q_n^{\zeta^{(o)}}$, there should be a natural way to go back and forth between them.

Proposition 2. *Suppose that ζ satisfies Condition 1 for $* = (o)$. Define $\hat{\zeta}$ by $\hat{\zeta}_n = n!\zeta_n$. Suppose that T is distributed like $Q_n^{\zeta^{(o)}}$ and let U be a uniformly random ordering of $\{1, \dots, n\}$ independent of T . Define $\hat{T} \in \mathcal{T}_n$ to be the tree obtained from T by labeling the leaves of T by U and forgetting the ordering of T . Then $\hat{T}$ is distributed like Q_n^ζ .*

Results of this type connecting plane and labeled trees where the size of a tree is given by the number of its vertices can be traced back to [10, 13, 14]. See [15] for a more complete history. Our proposition is analogous to an implicit discussion in [1, 2] as well as Theorem 7.1 in [15], which considered the case where the size of a tree is given by the number of its vertices. To prove this proposition, we will need some notation. For a rooted ordered tree x let $\text{shape}(x)$ be the rooted unordered tree obtained by forgetting the order on x . Similarly, for $t \in \mathcal{T}$, $\text{shape}(t)$ is defined to be the rooted unlabeled tree obtained from forgetting the labeling of t . For $t \in \mathcal{T}$, $x \in \mathcal{T}_n^{(o)}$, and a rooted unordered tree y define $\#\text{labels}_t(x)$ to be the number of ways to label the leaves of x such that when you forget the order on x you get t and $\#\text{ordered}(y)$ to be the number of ordered trees whose shape is y . Observe that $\#\text{labels}_t(x)$ depends only on $\text{shape}(x)$, so we will abuse our notation and write $\#\text{labels}_t(\text{shape}(x))$.

Proof. Observe that

$$\mathbb{P}(\hat{T} = t) = \sum_{x \in \mathcal{T}_n^{(o)}} \mathbb{P}(T = x) \mathbb{P}(\hat{T} = t | T = x).$$

Furthermore, observe that

$$\mathbb{P}(\hat{T} = t | T = x) = \frac{\#\text{labels}_t(\text{shape}(x))}{n!},$$

Observe that $\#\text{labels}_t(\text{shape}(x)) = 0$ unless $\text{shape}(t) = \text{shape}(x)$. Furthermore, $\mathbb{P}(T = x)$ depends only on $\text{shape}(x)$, and is given by

$$\mathbb{P}(T = x) = \frac{\prod_{v \in \text{shape}(x)} \zeta_{\deg(v)}}{\sum_{s \in \mathcal{T}_n^{(o)}} w_\zeta(s)}.$$

Consequently we have

$$(2.2) \quad \mathbb{P}(\hat{T} = t) = \frac{\#\text{ordered}(\text{shape}(t)) \prod_{v \in \text{shape}(t)} \zeta_{\deg(v)} \frac{\#\text{labels}_t(\text{shape}(t))}{n!}}{\sum_{s \in \mathcal{T}_n^{(o)}} w_\zeta(s)}.$$

But

$$(2.3) \quad \#\text{ordered}(\text{shape}(t)) \#\text{labels}_t(\text{shape}(t)) = \prod_{v \in \text{shape}(t)} (\deg(v)!).$$

This is because both sides count the number of distinct leaf-labeled ordered trees that equal t upon forgetting their order. On the left hand side, you pick a ordered tree and the label it and, on the right hand side, you label an unordered tree with the appropriate shape and then order the children of each vertex.

Therefore we have

$$\mathbb{P}(\hat{T} = t) = \frac{w_{\hat{\zeta}}(t)}{n! \sum_{s \in \mathcal{T}_n^{(o)}} w_\zeta(s)}.$$

The last step is to observe that

$$n! \sum_{s \in \mathcal{T}_n^{(o)}} w_\zeta(s) = \sum_{s \in \mathcal{T}_n} w_{\hat{\zeta}}(s).$$

This is because for $s \in \mathcal{T}_n^{(o)}$, there are $n!$ rooted ordered leaf-labeled trees whose ordered tree is s upon forgetting the labeling, so the left hand side is the weighted number of rooted ordered leaf-labeled trees with n leaves. Furthermore, we have already noted above that for $s \in \mathcal{T}_n$, there are $\prod_{v \in s} (\deg(v)!)^n$ rooted ordered leaf-labeled trees whose labeled tree is s upon forgetting the ordering. Thus the right hand side is also the weighted number of rooted ordered leaf-labeled trees with n leaves. Note that this step also shows that $\hat{\zeta}$ satisfies Condition 1 for $*$ being nothing. $\square$

Combining with tilting, we have the following corollary.

Corollary 2. *Let ζ satisfy Condition 1 with $*$ being nothing and $\zeta_0 = 1$. Suppose there exist $r > 0$ and $s > 0$ satisfying $s = r + G_\xi(s)$ and $G'_\xi(s) \leq 1$. Define $\xi = (\xi_i)_{i=0}^\infty$ by $\xi_0 = rs^{-1}$ and $\xi_j = s^{j-1}\zeta_j/j!$ for $j \geq 1$. Note that ξ is a probability distribution on $\mathbb{Z}_+$. Let T be a Galton-Watson tree with offspring distribution ξ and construct $\hat{T}$ by labeling the leaves of T uniformly at random with $\{1, \dots, |T|\}$, independently of T . Then $\mathbb{P}(\hat{T} \in \cdot | |T| = n) = Q_n^\zeta(\cdot)$ for all $n \geq 1$ such that Q_n^ζ is defined. Furthermore, for n such that Q_n^ζ is not defined, $P(|T| = n) = 0$.*

2.2. Schröder's problems. In this section we record which of the trees above correspond to the trees that appear in Schröder's problems. The proofs of the claims here are simple applications of the results in Section 2.1.

The first problem: The trees here are uniform binary rooted ordered unlabeled trees. We can obtain these by taking $*$ = (o) and $\zeta_0 = \zeta_2 = 1$ and $\zeta_i = 0$ for $i \notin \{0, 2\}$. Letting ξ be the probability distribution given by $\xi_0 = \xi_2 = 1/2$ and T be a Galton-Watson tree with offspring distribution ξ , we have that T conditioned to have n leaves is a uniform binary rooted ordered unlabeled tree with n leaves. Also note that T is critical and the variance of ξ is equal to one.

The second problem: These are uniform rooted ordered trees with no vertices of out degree one. These were dealt with in Corollary 1

The third problem: These are uniform binary unordered leaf-labeled trees. We can obtain these by taking $*$ to be nothing and $\zeta_0 = \zeta_2 = 1$ and $\zeta_i = 0$ for $i \notin \{0, 2\}$. In this case, if T is the Galton-Watson tree defined in the first problem and $\hat{T}$ is defined as in Corollary 2, then $\hat{T}$ conditioned to have n leaves is a uniform binary unordered leaf-labeled tree with n leaves.

The fourth problem: These are uniform rooted unordered leaf-labeled trees with no vertices with out-degree 1. We can obtain these by taking $*$ to be nothing and $\zeta_1 = 0$ and $\zeta_i = 1$ for $i \neq 1$. We define a probability distribution ξ by

$$\xi_0 = \frac{2 \log(2) - 1}{\log(2)}, \quad \xi_1 = 0, \quad \text{and } \xi_j = \frac{(\log(2))^{j-1}}{j!} \text{ for } j \geq 2.$$

Note that ξ has mean 1 and variance $\text{var}(\xi) = 2 \log 2$. Letting T be a Galton-Watson tree with offspring distribution ξ and defining $\hat{T}$ is as in Corollary 2, we have that $\hat{T}$ conditioned to have n leaves is a uniform unordered leaf-labeled tree with no vertices of out degree one and n leaves.

2.3. Gibbs trees. Above we saw a natural way to put probability measures on $\mathcal{T}_n$ that are concentrated on fragmentation trees (the trees appearing in Schröder's fourth problem); namely, take $\zeta_1 = 0$. Another natural type of probability to put on fragmentation trees is a *Gibbs model*, which we now describe. First, we need to set up the natural framework in which to view fragmentation trees. The idea is that, while in Schröder's fourth problem we have an arbitrary set bracketing, for fragmentations we recursively partition a set. This dynamic view of constructing a set bracketing makes Gibbs models quite natural.

Definition 1 ([12]). *A fragmentation of the finite set B is a collection $\mathbf{t}_B$ of non-empty subsets of B such that*

- (1) $B \in \mathbf{t}_B$
- (2) *If $\#B \geq 2$ then there is a partition of B into $k \geq 2$ parts $B_1, \dots, B_k$, called the children of B , such that*

$$\mathbf{t}_B = \{B\} \cup \mathbf{t}_{B_1} \cup \dots \cup \mathbf{t}_{B_k},$$

where $\mathbf{t}_{B_i}$ is a fragmentation of B_i .

We can naturally consider $\mathbf{t}_B$ as a tree whose vertices are the elements of $\mathbf{t}_B$ and whose edges are defined by the parent child relationship. Considering the properties of such a tree leads naturally to the following definition of a *fragmentation tree* on B .

Definition 2. *A fragmentation tree T on n leaves is a rooted tree such that*

- (1) *The root of T does not have degree 1,*
- (2) *T has no non-root vertices of degree 2,*
- (3) *The leaves of T are labeled by a set B with $\#B = n$. We denote the label of a leaf v by $\ell(v)$.*

The idea of the Gibbs model is that, at each step in the fragmentation the next step is distributed according to multiplicative weights depending on the block sizes. We first take a sequence $\{\alpha_n\}$, $\alpha_n \geq 0$ of weights and a Gibbs weight, which is a function $g : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$ with $g(0) = 0$. Then, for $n \geq 2$, define a normalization constant

$$Z(n) = \sum_{\{B_1, \dots, B_k\}} \alpha_k \prod_{j=1}^k g(\#B_j),$$

where the sum is over unordered partitions of $[n]$ into at least two elements. Whenever we write a formula like this, we assume that each block B_i is nonempty. Now, define the probability of a partition of $[n]$ by

$$P_n^{g, \alpha}(B_1, \dots, B_k) = p(\#B_1, \dots, \#B_k) = \frac{\alpha_k \prod_{j=1}^k g(\#B_j)}{Z(n)}.$$

The probability of a fragmentation X of $[n]$ is then defined as

$$P_n^{g, \alpha}(X) = \prod_{B \in X} P(B_1, \dots, B_k),$$

where $B_1, \dots, B_k$ are the children of B . Using the correspondence between fragmentations and fragmentation trees, for $T_n \in \mathcal{T}_n$, we define $P_n^{g, \alpha}(T_n)$ to be $P_n^{g, \alpha}(X)$ where X is the

fragmentation determined by T_n . The probabilistic properties of Gibbs models are studied in [12].

Theorem 2. *Suppose that ζ satisfies Condition 1 with $*$ being nothing and $\zeta_1 = 0$. Define $\alpha_k = \zeta_k$ and $g(k) = k![z^k]C_\zeta(z)$. Then $Z(n) = g(n)$ and $Q_n^\zeta = P_n^{g,\alpha}$. Furthermore, given a nonnegative weight sequence α and a Gibbs weight g such that $Z(n) = g(n)$, there is a ζ satisfying Condition 1 with $*$ being nothing and $\zeta_1 = 0$ such that $Q_n^\zeta = P_n^{g,\alpha}$.*

Proof. Arguing similarly as in (2.1), we see that for $n \geq 2$

$$Z(n) = \sum_{\{B_1, \dots, B_k\}} \alpha_k \prod_{j=1}^k g(\#B_j) = \sum_{k=2}^{\infty} \frac{\alpha_k}{k!} \sum_{\substack{(n_1, \dots, n_k) \in \mathbb{N}^k \\ n_1 + \dots + n_k = n}} \binom{n}{n_1, \dots, n_k} \prod_{j=1}^k g(n_j) = n![z^n]C_\zeta(z).$$

Consequently we have $Z(n) = g(n)$. Using this, one proves inductively that $P_n^{g,\alpha}(T_n) = Q_n^\zeta(T_n)$. Furthermore, observe that the condition $Z(n) = g(n)$ implies that there is a weight sequence $(\zeta_i)_{i \geq 0}$ from which the fragmentation model can be derived in the above manner; just take $\zeta_0 = g(1)$, $\zeta_1 = 0$, and $\zeta_k = \alpha_k$ for $k \geq 2$. $\square$

When we have $Z(n) = g(n)$, the model is called a *combinatorial* Gibbs model. This is justified by the fact that, in this case, $Z(n)$ (and thus $g(n)$) is the weighted number of trees with n leaves. For example, if we let $g(n)$ be the number of fragmentation trees with n leaves, and $\alpha_k = 1$ for $k \geq 2$, we then see that

$$Z(n) = \sum_{\{B_1, \dots, B_k\}} \prod_{j=1}^k g(\#B_j).$$

The right hand side of this equation is just the sum over partitions at the root of a fragmentation tree with n leaves of the number of fragmentation trees with that partition at the root, which is precisely the number of fragmentation trees with n leaves. That is, $Z(n) = g(n)$.

Note that combinatorial Gibbs models are a generalization of the hierarchies studied in [7] and, as previously observed, a special case of the Gibbs models introduced in [12].

3. SCALING LIMITS

We now turn to scaling limits of the models of trees we have been discussing. Fortunately for us, the heavy lifting has already been done in [16]. In order to use the results from that paper, we must first introduce the formalism required to handle limits of random metric measure spaces.

3.1. Trees as metric measure spaces. The trees we have been talking about can naturally be considered as metric spaces with the graph metric. That is, the distance between two vertices is the number of edges on the path connecting them. Let (t, d) be a tree equipped with the graph metric. For $a > 0$, we define at to be the metric space (t, ad) , i.e. the metric is scaled by a . This is equivalent to saying the edges have length a rather than length 1 in the definition of the graph metric. More, generally we can attach a positive length to each edge in t and use these to in the definition of the graph metric. Moreover, the trees we are dealing with are rooted so we consider (t, d) as a pointed metric space with the root as the

point. Moreover, we are concerned with the leaves, so we attach a measure μ_t , which is the uniform probability measure on the leaves of t . If we have a random tree T , this gives rise to a random pointed metric measure space $(T, d, \text{root}, \mu_T)$. To make this last concept rigorous, we need to put a topology on pointed metric measure spaces. This is hard to do in general, but note that the pointed metric measure spaces that come from the trees we are discussing are compact.

Let $\mathcal{M}_w$ be the set of equivalence classes of compact pointed metric measure spaces (equivalence here being up to point and measure preserving isometry). It is worth pointing out that $\mathcal{M}_w$ actually is a set in the sense of ZFC, though this takes some work to show. We metrize $\mathcal{M}_w$ with the pointed Gromov-Hausdorff-Prokhorov metric (see [8]). Fix $(X, d, \rho, \mu), (X', d', \rho, \mu') \in \mathcal{M}_w$ and define

$$d_{\text{GHP}}(X, X') = \inf_{(M, \delta)} \inf_{\substack{\phi: X \rightarrow M \\ \phi': X' \rightarrow M}} [\delta(\phi(\rho), \phi'(\rho')) \vee \delta_H(\phi(X), \phi'(X')) \vee \delta_P(\phi_*\mu, \phi'_*\mu')],$$

where the first infimum is over metric spaces (M, δ) , the second infimum is over isometric embeddings ϕ and ϕ' of X and X' into M , δ_H is the Hausdorff distance on compact subsets of M , and δ_P is the Prokhorov distance between the pushforward $\phi_*\mu$ of μ by ϕ and the pushforward $\phi'_*\mu'$ of μ' by ϕ' . Again, the definition of this metric has potential to run into set-theoretic difficulties, but they are not terribly difficult to resolve.

Proposition 3 (Proposition 1 in [8]). *The space $(\mathcal{M}_w, d_{\text{GHP}})$ is a complete separable metric space.*

An $\mathbb{R}$ -tree is a complete metric space (T, d) with the following properties:

- For $v, w \in T$, there exists a unique isometry $\phi_{v,w} : [0, d(v, w)]$ with $\phi_{v,w}(0) = v$ to $\phi_{v,w}(d(v, w)) = w$.
- For every continuous injective function $c : [0, 1] \rightarrow T$ such that $c(0) = v$ and $c(1) = w$, we have $c([0, 1]) = \phi_{v,w}([0, d(v, w)])$.

If (T, d) is an $\mathbb{R}$ -tree, every choice of root $\rho \in T$ and probability measure μ on T yields an element (T, d, ρ, μ) of $\mathcal{M}_w$. With this choice of root also comes a height function $\text{ht}(v) = d(v, \rho)$. The leaves of T can then be defined as a point $v \in T$ such that v is not in $[[\rho, w[[:= \phi_{\rho,w}([0, \text{ht}(w)))$ for any $w \in T$. The set of leaves is denoted $\mathcal{L}(T)$.

Definition 3. *A continuum tree is an $\mathbb{R}$ -tree (T, d, ρ, μ) with a choice of root and probability measure such that μ is non-atomic, $\mu(\mathcal{L}(T)) = 1$, and for every non-leaf vertex w , $\mu\{v \in T : [[\rho, v]] \cap [[\rho, w]] = [[\rho, w]]\} > 0$.*

The last condition says that there is a positive mass of leaves above every non-leaf vertex. We will usually just refer to a continuum tree T , leaving the metric, root, and measure as implicit. A continuum random tree (CRT) is an $(\mathcal{M}_w, d_{\text{GHP}})$ valued random variable that is almost surely a continuum tree.

3.2. The Brownian continuum random tree. Continuous functions give a nice way of constructing $\mathbb{R}$ -trees. Suppose that $f : [0, 1] \rightarrow \mathbb{R}_+$ is continuous and $f(0) = f(1) = 0$. We can define a pseudo-metric on $[0, 1]$ by $d_f(a, b) = f(a) + f(b) - 2 \min_{a \leq t \leq b} f(t)$ for $a \leq b$. Define an equivalence relation by $a \sim b$ if and only if $d_f(a, b) = 0$. Letting $T_f = [0, 1] / \sim$, we obtain a metric space (T_f, d_f) . Theorem 2.1 in [6] tells us that (T_f, d_f) is a compact $\mathbb{R}$ -tree.

Letting $\rho : [0, 1] \rightarrow T_f$ be the natural map taking a point to its equivalence class, we can take $\rho(0)$ as the root of T_f . We then obtain an element of $\mathcal{M}_w$ by equipping T_f with the measure μ_f induced by the pushforward of Lebesgue measure on $[0, 1]$ onto T_f by ρ .

Definition 4. *Let $(e(t), 0 \leq t \leq 1)$ be standard Brownian excursion. The Brownian continuum random tree, denoted T^{Br} , is continuum random tree $(T_e, d_e, \rho(0), \mu_e)$.*

Note that the elementary properties of Brownian excursion imply that T^{Br} actually is almost surely a continuum tree. It is also worth noting that in our formalism the Brownian continuum random tree originally defined by Aldous in [3] corresponds to $(T_{2e}, d_{2e}, \rho(0), \mu_{2e})$, but the convention has since shifted to the one we have adopted here (see e.g. [11]).

3.3. Convergence theorems. With this machinery, we can import the following special case of Theorem 1 in [16].

Theorem 3 (Theorem 1 in [16]). *Let T be a critical Galton-Watson tree with offspring distribution ξ such that $0 < \sigma^2 = \text{var}(\xi) < \infty$. Suppose that for sufficiently large n the probability that T has exactly n leaves is positive and for such n let T_n be T conditioned to have exactly n leaves, considered as a rooted unordered tree with edge lengths 1 and the uniform probability distribution μ_{T_n} on its leaves. Then*

$$\frac{1}{\sqrt{n}}T_n \xrightarrow{d} \frac{2}{\sigma\sqrt{\xi_0}}T^{Br},$$

where the convergence is with respect to the rooted Gromov-Hausdorff-Prokhorov topology.

Proof of Theorem 1. Corollary 2 provides for the direct conversion of the results of Theorem 3 into results about the trees in Schröder's problems. $\square$

4. EXPLICIT COMPUTATIONS USING ANALYTIC COMBINATORICS

The convergence result above is a powerful theorem for obtaining asymptotics of various tree statistics, but it is difficult to prove and, as a result, asymptotics thus obtained can seem mysterious. Consequently it is worth noting we can obtain a number of asymptotic results directly using analytic combinatorics. This analytic approach is based on considering the asymptotics of generating functions. The primary source for asymptotics in general is [7], which develops the theory with extensive examples.

Our main goal in this section is to develop the general framework of additive functionals for leaf-labeled trees whose size is counted by their number of leaves and use this to find the asymptotic distribution of the height of a uniformly randomly chosen leaf. We also find the limit of the expected height of a random leaf. These computations are meant to be illustrative and by no means exhaust the power of analytic combinatorics framework. Indeed, it seems that most of the techniques used to study for simple varieties of trees (see [7] for a summary of the extensive work in this area) have close analogs that will provide results about the trees we are considering here.

4.1. Analytic background. We summarize some of the fundamental results here, but make no attempt to prove them. The approach is based on the asymptotics of several universal functions. Recall that if $f(z)$ is either a formal power series, $[z^n]f(z)$ denotes the coefficient of z^n . Similarly, if $f: \mathbb{C} \rightarrow \mathbb{C}$ is analytic at 0 then $[z^n]f(z)$ denotes the coefficient of z^n in the power series expansion of f at 0.

Proposition 4. *Let $f(z) = (1 - z)^{1/2}$, $g(z) = (1 - z)^{-1/2}$, and $h(z) = (1 - z)^{-1}$. Then $[z^n]f(z) \sim -1/2\sqrt{\pi n^3}$, $[z^n]g(z) \sim 1/\sqrt{n\pi}$, and $[z^n]h(z) = 1$.*

To use these classical results we need a special type of analyticity called Δ -analyticity, which we now define.

Definition 5 (Definition VI.I p. 389 [7]). *Given two number ϕ and R with $R > 1$ and $0 < \phi < \pi/2$, the open domain $\Delta(\phi, R)$ is defined as*

$$\Delta(\phi, R) = \{z \mid |z| < R, z \neq 1, |\arg(z - 1)| > \phi\}.$$

A for a complex number ζ a domain D is a Δ -domain at ζ if there exist ϕ and R such that $D = \zeta\Delta(\phi, R)$. A function is Δ -analytic if it is analytic on a Δ -domain.

Let

$$\mathcal{S} = \{(1 - z)^{-\alpha} \lambda(z)^\beta \mid \alpha, \beta \in \mathbb{C}\} \quad \lambda(z) \equiv \frac{1}{z} \log \frac{1}{1 - z}.$$

Theorem 4 (Theorem VI.4 p. 393 [7]). *Let $f(z)$ be a function analytic at 0 with a singularity at ζ , such that $f(z)$ can be continued to a domain of the form $\zeta\Delta_0$, for a Δ -domain Δ_0 . Assume that there exist two function σ and τ , where σ is a (finite) linear combination of elements of $\mathcal{S}$ and $\tau \in \mathcal{S}$, so that*

$$f(z) = \sigma(z/\zeta) + O(\tau(z/\zeta)) \quad \text{as } z \rightarrow \zeta \text{ in } \zeta\Delta_0.$$

Then the coefficients of $f(z)$ satisfy the asymptotic estimate

$$f_n = \zeta^{-n} \sigma_n + O(\zeta^{-n} \tau_n^*),$$

where $\tau_n^* = n^{a-1}(\log n)^b$, if $\tau(z) = (1 - z)^{-a} \lambda(z)^b$.

Occasionally we will also need to deal with derivatives and the next theorem shows us how this is done.

Theorem 5 (Theorem VI.8 p. 419 [7]). *Let $f(z)$ be Δ -analytic with singular expansion near its singularity of the simple form*

$$f(z) = \sum_{j=0}^J c_j (1 - z)^{a_j} + O((1 - z)^A).$$

Then, for each integer $r > 0$, the derivative $f^{(r)}(z)$ is Δ -analytic. The expansion of the derivative at its singularity is obtained through term by term differentiation:

$$\frac{d^r}{dz^r} f(z) = (-1)^r \sum_{j=0}^J c_j \frac{\Gamma(a_j + 1)}{\Gamma(a_j + 1 - r)} (1 - z)^{a_j - r} + O((1 - z)^{A-r}).$$

The generating functions we will work with fall into the smooth implicit-function schema, which provides a way to derive coefficient asymptotics from functional equations.

Definition 6 (Definition VII.4 p. 467 [7]). *Let $y(z)$ be a function analytic at 0, $y(z) = \sum_{n \geq 0} y_n z^n$, with $y_0 = 0$ and $y_n \geq 0$. The function is said to belong to the smooth implicit-function schema if there exists a bivariate function $G(z, w)$ such that*

$$y(z) = G(z, y(z)),$$

where $G(z, w)$ satisfies the following conditions.

- (i) $G(z, w) = \sum_{m, n \geq 0} g_{m, n} z^m w^n$ is analytic in a domain $|z| < R$ and $|w| < S$, for some $R, S > 0$.
- (ii) The coefficients of G satisfy $g_{m, n} \geq 0$, $g_{0, 0} = 0$, $g_{0, 1} \neq 1$, and $g_{m, n} > 0$ for some m and for some $n \geq 2$.
- (iii) There exist two numbers r and s such that $0 < r < R$ and $0 < s < S$, satisfying the system of equations

$$G(r, s) = s, \quad G_w(r, s) = 1, \quad \text{with } r < R, \quad s < S,$$

which is called the characteristic system.

Definition 7 (Definition IV.5 p. 266 [7]). *Consider the formal power series $f(z) = \sum f_n z^n$. The series f is said to admit span d if for some r*

$$\{f_n\}_{n=0}^{\infty} \subseteq r + d\mathbb{Z}_+.$$

The largest span is the period of f . If f has period 1, then f is aperiodic.

With this definition, we get the following theorem. It is worth noting that this result appears in several places in the literature. We give the version that appears as Theorem VII.3 on page 468 of [7]. In that source it is footnoted that many statements occurring previously in the literature contained errors, so caution is advised.

Theorem 6 (Theorem VII.3 p. 468 [7]). *Let $y(z)$ belong to the smooth implicit-function schema defined by $G(z, w)$, with (r, s) the positive solution of the characteristic system. Then $y(z)$ converges at $z = r$, where it has a square root singularity,*

$$y(z) \underset{z \rightarrow r}{=} s - \gamma \sqrt{1 - z/r} + O(1 - z/r), \quad \gamma \equiv \sqrt{\frac{2rG_z(r, s)}{G_{ww}(r, s)}},$$

the expansion being valid in a Δ -domain. If, in addition, $y(z)$ is aperiodic, then r is the unique dominant singularity of y and the coefficients satisfy

$$[z^n]y(z) \underset{n \rightarrow \infty}{=} \frac{\gamma}{2\sqrt{\pi n^3}} r^{-n} (1 + O(n^{-1})).$$

We will also need the following theorem.

Theorem 7 ((A special case of) Theorem IX.16 p. 709 [7]). *Let $H(z)$ be Δ -continuable and of the form $H(z) = \sigma - h(1 - z/\rho)^{1/2} + O(1 - z/\rho)$ and let $k_n = x_n n^{1/2}$ for x_n in any compact subinterval of $(0, \infty)$. Then*

$$[z^n]H(z)^{k_n} \sim \sigma^{k_n} \rho^{-n} \frac{h k_n}{2\sigma \sqrt{\pi n^3}} \exp\left(-\frac{h^2 k_n^2}{4\sigma^2 n}\right).$$

4.2. Restricting the generality. So far we have been considering a very general situation. However, in what follows we will be doing computations that are tedious to do in full generality. Consequently, we will restrict the generality. In particular, we will let $\zeta = (\zeta_i)_{i \geq 0}$ be a sequence of non-negative weights such that $\zeta_0 = 1$, $\zeta_1 = 0$, $\gcd\{k : \zeta_k \neq 0\} = 1$, and

$$G_\zeta(z) = \sum_{j=2}^{\infty} \zeta_j \frac{z^j}{j!}$$

is entire. These conditions can be relaxed, but doing so makes the analysis more difficult.

Proposition 5. *With ζ as above, C_ζ (defined in Section 2.1) belongs to the smooth implicit-function schema with $G(z, w) = z + G_\zeta(w)$. Furthermore, in the case where ζ corresponds to Schröder's third problem $(r, s) = (1/2, 1)$ and in the case of the fourth problem, we have $(r, s) = (2 \log(2) - 1, \log(2))$. Additionally*

$$[z^n]C_\zeta(z) \sim \frac{\gamma}{2\sqrt{\pi n^3}} r^{-n}, \quad \gamma = \sqrt{\frac{2r}{G_\zeta''(s)}}.$$

Proof. All that really needs to be checked is that the characteristic system has a positive solution. For $G(z, w) = z + G_\zeta(w)$, the characteristic system is $s = r + G_\zeta(s)$ and $G_\zeta'(s) = 1$. Using that G_ζ is entire, $G_\zeta'(0) = 0$ and $G_\zeta'(+\infty) = +\infty$, and G_ζ' is increasing on $\mathbb{R}_+$, the intermediate value theorem yields $s > 0$. An easy computation yields that $G_\zeta(s) < sG_\zeta'(s) = s$, so $r > 0$ as well. $\square$

4.3. The height of a random leaf. Let H_n be the height of a randomly chosen leaf from a tree in $\mathcal{T}_n$. Specifically, to get H_n , we choose a tree T_n from $\mathcal{T}_n$ according to P_n and then choose a leaf uniformly at random from T_n . Our main result in this section is the following theorem.

Theorem 8.

$$\frac{\lambda}{\sqrt{n}} H_n \xrightarrow{d} \text{Rayleigh}(1),$$

for $\lambda = \sqrt{G_\zeta''(s)r}$. In Schröder's third problem $\lambda = 1/\sqrt{2}$, and in the fourth problem $\lambda = \sqrt{4 \log(2) - 2}$.

Our approach will be that of additive functionals, whose theory we now develop. We parallel the development of these functions in [7], p. 457. Their work was done for simple varieties of trees whose size was determined by the number of vertices. Here we work with trees whose size is determined by the number of leaves.

For a rooted unordered tree t whose leaves are labeled by $B \subseteq \mathbb{N}$, let $\tilde{t} \in \mathcal{T}$ be the tree that results from relabeling the leaves of t by the unique increasing bijection from B to $\{1, 2, \dots, |B|\}$ (where $|B|$ is the cardinality of B). Suppose we have functions $\xi, \theta, \psi : \mathcal{T} \rightarrow \mathbb{R}$ satisfying the relation

$$\xi(t) = \theta(t) + \sum_{j=1}^{\deg(t)} \psi(\tilde{t}_j),$$

where $\deg(t)$ is the root degree of t and the $\{t_j\}$ are the root subtrees of t ordered in increasing order of the leaf with the smallest label. Letting $\bullet$ denote the tree on one leaf, we note that $\deg(\bullet) = 0$, so in particular $\xi(\bullet) = \theta(\bullet)$. Define the exponential generating functions

$$\Xi(z) = \sum_{t \in \mathcal{T}} \xi(t) w(t) \frac{z^{|t|}}{|t|!}, \quad \Theta(z) = \sum_{t \in \mathcal{T}} \theta(t) w(t) \frac{z^{|t|}}{|t|!}, \quad \text{and} \quad \Psi(z) = \sum_{t \in \mathcal{T}} \psi(t) w(t) \frac{z^{|t|}}{|t|!}.$$

Our results make use of the following lemma, which is a relation of formal power series.

Lemma 1. *Let $C_\zeta(z)$ be the exponential generating function for $\mathcal{C}$. Then*

$$(4.1) \quad \Xi(z) = \Theta(z) + G'_\zeta(C_\zeta(z))\Psi(z).$$

In the purely recursive case where $\xi \equiv \psi$ we have

$$(4.2) \quad \Xi(z) = \frac{\Theta(z)}{1 - G'_\zeta(C_\zeta(z))} = C'_\zeta(z)\Theta(z).$$

Proof. We clearly have

$$\Xi(z) = \Theta(z) + \tilde{\Psi}(z), \quad \text{where} \quad \tilde{\Psi}(z) = \sum_{t \in \mathcal{T}} \left(w(t) \frac{z^{|t|}}{|t|!} \sum_{j=1}^{\deg(t)} \psi(\tilde{t}_j) \right).$$

Decomposing by root degree and using that $\xi(\bullet) = \theta(\bullet)$, we have

$$\begin{aligned} \tilde{\Psi}(z) &= \sum_{r \geq 1} \sum_{\deg(t)=r} \zeta_r \prod_{j=1}^r w(\tilde{t}_j) \frac{z^{|t_1|+\dots+|t_r|}}{(|t_1|+\dots+|t_r|)!} (\psi(\tilde{t}_1) + \dots + \psi(\tilde{t}_r)) \\ &= \sum_{r \geq 1} \left[\frac{\zeta_r}{r!} \sum_{(t_1, \dots, t_r) \in \mathcal{T}^r} \prod_{j=1}^r w(t_j) \binom{|t_1|+\dots+|t_r|}{|t_1|, \dots, |t_r|} \frac{z^{|t_1|+\dots+|t_r|}}{(|t_1|+\dots+|t_r|)!} (\psi(t_1) + \dots + \psi(t_r)) \right] \\ &= \sum_{r \geq 1} \left[\frac{\zeta_r}{r!} \sum_{(t_1, \dots, t_r) \in \mathcal{T}^r} \prod_{j=1}^r w(t_j) \frac{z^{|t_1|+\dots+|t_r|}}{|t_1|! \dots |t_r|!} (\psi(t_1) + \dots + \psi(t_r)) \right] \\ &= \sum_{r \geq 1} \frac{\zeta_r}{(r-1)!} C_\zeta(z)^{r-1} \Psi(z) \\ &= G'_\zeta(C_\zeta(z))\Psi(z). \end{aligned}$$

This yields (4.1). In the recursive case, we have $\Xi(z) = \Theta(z) + G'_\zeta(C_\zeta(z))\Xi(z)$. Solving for $\Xi(z)$ gives the first equality in (4.2). To get the second, we differentiate (2.1) to get $C'_\zeta(z) = 1 + G'_\zeta(C_\zeta(z))C'_\zeta(z)$. Solving for $C'_\zeta(z)$ gives $C'_\zeta(z) = 1/(1 - G'_\zeta(C_\zeta(z)))$, from which the second equality in (4.2) is immediate. $\square$

Two immediate applications are to counting the weighted numbers of leaves and vertices of a given height.

Theorem 9. *The expected number of leaves at height k converges to $G''_\zeta(s)rk$ and the expected number of nodes at height k converges to $sG''_\zeta(s)k + 1$.*

Proof. Let $\xi_k(t)$ be the number of leaves of height k in t , so that $\xi_k(t)w(t)$ is the weighted number of leaves of height k . Define $\Xi_k = \sum_t \xi_k(t)w(t)z^{|t|}/|t|!$. For $k \geq 1$ we apply the lemma with $\xi = \xi_k$, $\theta = 0$ and $\psi = \xi_{k-1}$ to obtain

$$\Xi_k(z) = G'_\zeta(C_\zeta(z))\Xi_{k-1}(z),$$

which easily yields

$$\Xi_k(z) = [G'_\zeta(C_\zeta(z))]^k \Xi_0(z) = z [G'_\zeta(C_\zeta(z))]^k.$$

Letting $\Lambda_k(z)$ be the generating function for the weighted number of vertices of height k we similarly get

$$\Lambda_k(z) = [G'_\zeta(C_\zeta(z))]^k \Lambda_0(z) = C_\zeta(z) [G'_\zeta(C_\zeta(z))]^k.$$

Using these forms, we are able to compute asymptotics. Expanding G'_ζ about s , we have that $G'_\zeta(z) = 1 + G''_\zeta(s)(z - s) + O((z - s)^2)$. Plugging in the asymptotic expansion of C_ζ we get from Proposition 5 and Theorem 6 and doing some algebra, we have

$$(4.3) \quad G'_\zeta(C_\zeta(z)) = 1 - G''_\zeta(s)\gamma\sqrt{1 - z/r} + O(1 - z/r).$$

Hence, using that $(1 - z)^k = 1 - kz + O(z^2)$, we see that

$$[G'_\zeta(C_\zeta(z))]^k = 1 - G''_\zeta(s)k\gamma\sqrt{1 - z/r} + O(1 - z/r).$$

Thus, using Theorem 4, we have

$$[z^n]\Xi_k(z) = [z^n]z [G'_\zeta(C_\zeta(z))]^k \sim \frac{G''_\zeta(s)k\gamma}{2r^{n-1}\sqrt{\pi n^3}},$$

and, similarly with a bit more algebra,

$$[z^n]\Lambda_k(z) = [z^n]C_\zeta(z) [G'_\zeta(C_\zeta(z))]^k \sim \frac{\gamma(skG''_\zeta(s) + 1)}{2r^n\sqrt{\pi n^3}}.$$

Using the result on p. 474 of [7], that $[z^n]C_\zeta(z) \sim \gamma/2r^n\sqrt{\pi n^3}$, we find that

$$E_{\mathcal{T}_n}(\xi_k) = \frac{n![z^n]\Xi_k(z)}{n![z^n]C_\zeta(z)} \sim G''_\zeta(s)rk.$$

Letting $\zeta_k : \mathcal{T} \rightarrow \mathbb{Z}$ be the number of nodes of height k in t , we have that

$$E_{\mathcal{T}_n}(\zeta_k) = \frac{n![z^n]\Lambda_k(z)}{n![z^n]C_\zeta(z)} \sim sG''_\zeta(s)k + 1.$$

□

The proof of Theorem 8 is similar, but we make use of Theorem 7 for the asymptotics.

Proof of Theorem 8. Let $\{k_n\}$ be a sequence of integers varying such that $ck_n/n^{1/2} \rightarrow x \in (0, \infty)$ for some $c > 0$. By Theorem 7 and equation (4.3) we see that

$$(4.4) \quad [z^n][G'_\zeta(C_\zeta(z))]^{k_n} \sim r^{-n} \frac{G''_\zeta(s)\gamma}{2\sqrt{\pi n^3}} k_n \exp\left(-\frac{G''_\zeta(s)^2\gamma^2 k_n^2}{4n}\right).$$

Therefore

$$[z^n]\Xi_{k_n}(z) \sim r^{-(n-1)} \frac{G''_\zeta(s)\gamma}{2\sqrt{\pi n^3}} k_{n-1} \exp\left(-\frac{G''_\zeta(s)^2 \gamma^2 k_{n-1}^2}{4(n-1)}\right).$$

This yields

$$\begin{aligned} E_{\mathcal{T}_n}(\xi_{k_n}) &\sim G''_\zeta(s) r k_{n-1} \exp\left(-\frac{G''_\zeta(s)^2 \gamma^2 k_{n-1}^2}{4(n-1)}\right) \\ &= G''_\zeta(s) r k_{n-1} \exp\left(-\frac{G''_\zeta(s) r k_{n-1}^2}{2(n-1)}\right). \end{aligned}$$

Note that $P(H_n = k) = E_{\mathcal{T}_n}(\xi_k)/n$. Observe that $\{k_n\}$ satisfies the hypotheses of the above theorem. Consequently, we have that

$$\begin{aligned} \frac{\sqrt{n}}{c} P_n\left(\frac{c}{\sqrt{n}} H_n = \frac{c}{\sqrt{n}} k_n\right) &= \frac{\sqrt{n}}{c} P(H_n = k_n) \\ &= \frac{1}{c\sqrt{n}} E_{\mathcal{T}_n}(\xi_{k_n}) \\ &\sim \frac{1}{c^2} G''_\zeta(s) r \frac{c k_{n-1}}{\sqrt{n}} \exp\left(-\frac{G''_\zeta(s) r}{2c^2} \frac{c^2 k_{n-1}^2}{(n-1)}\right) \\ &\rightarrow \frac{G''_\zeta(s) r}{c^2} x \exp\left(-\frac{G''_\zeta(s) r}{2c^2} x^2\right). \end{aligned}$$

The proof is finished by an application of a standard corollary of Scheffé's theorem (see Theorem 3.3 in [4] for an idea of the proof, just adapted for a distribution on $(0, \infty)$) and choosing $c = \sqrt{G''_\zeta(s)r}$. $\square$

In addition to proving convergence in distribution we can prove convergence of the first moment.

Theorem 10. $E_{\mathcal{T}_n} H_n \sim \sqrt{\frac{\pi}{2rG''_\zeta(s)}} n^{1/2}$.

The approach is to first compute the expected sum of the heights of the leaves of a tree.

Theorem 11. Let $\phi(t)$ be the sum of the heights of the leaves of t . Then $E_{\mathcal{T}_n} \phi \sim \sqrt{\frac{\pi}{2rG''_\zeta(s)}} n^{3/2}$.

Proof. Observe that

$$\phi(t) = |t| + \sum_{j=1}^{\deg(T)} \phi(\tilde{t}_j).$$

Let $\Phi(z)$ be the exponential generating function associated with ϕ . Applying Lemma 1, we have

$$\Phi(z) = z(C'_\zeta(z))^2.$$

By Theorem 5 we have

$$C'_\zeta(z) = \frac{\gamma}{2r} (1 - z/r)^{-1/2} + O(1).$$

Consequently,

$$(C'_\zeta(z))^2 = \frac{\gamma^2}{4r^2}(1 - z/r)^{-1} + ((1 - z/r)^{-1/2}).$$

Therefore

$$E_{\mathcal{T}_n} \phi = \frac{n![z^{n-1}](C'_\zeta(z))^2}{n![z^n]C_\zeta(z)} \sim \frac{\frac{\gamma^2 r^{-n+1}}{4r^2}}{\frac{\gamma}{2\sqrt{\pi n^3}} r^{-n}} = \frac{\gamma\sqrt{\pi}}{2r} n^{3/2} = \sqrt{\frac{\pi}{2rG''_\zeta(s)}} n^{3/2}.$$

□

Proof of Theorem 10. Simply observe that $E_{\mathcal{T}_n} H_n = \frac{1}{n} E_{\mathcal{T}_n} \phi$.

□

REFERENCES

- [1] David Aldous. The continuum random tree. I. *Ann. Probab.*, 19(1):1–28, 1991.
- [2] David Aldous. The continuum random tree. II. An overview. In *Stochastic analysis (Durham, 1990)*, volume 167 of *London Math. Soc. Lecture Note Ser.*, pages 23–70. Cambridge Univ. Press, Cambridge, 1991.
- [3] David Aldous. The continuum random tree. III. *Ann. Probab.*, 21(1):248–289, 1993.
- [4] Patrick Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons Inc., New York, second edition, 1999. A Wiley-Interscience Publication.
- [5] Michael Drmota. *Random trees*. SpringerWienNewYork, Vienna, 2009. An interplay between combinatorics and probability.
- [6] Thomas Duquesne and Jean-François Le Gall. Probabilistic and fractal aspects of Lévy trees. *Probab. Theory Related Fields*, 131(4):553–603, 2005.
- [7] Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. Cambridge University Press, Cambridge, 2009.
- [8] Bénédicte Haas and Grégory Miermont. Scaling limits of Markov branching trees, with applications to Galton-Watson and random unordered trees. *arXiv:1003.3632v2*, 2010.
- [9] Bénédicte Haas, Grégory Miermont, Jim Pitman, and Matthias Winkel. Continuum tree asymptotics of discrete fragmentations and applications to phylogenetic models. *Ann. Probab.*, 36(5):1790–1837, 2008.
- [10] V. F. Kolchin. Branching processes, random trees, and a generalized scheme of arrangements of particles. *Mathematical Notes*, 21:386–394, 1977. 10.1007/BF01788236.
- [11] Jean-François Le Gall. Random real trees. *Ann. Fac. Sci. Toulouse Math. (6)*, 15(1):35–62, 2006.
- [12] Peter McCullagh, Jim Pitman, and Mathias Winkel. Gibbs fragmentation trees. *Bernoulli*, 14:988–1002, 2008.
- [13] Yu. L. Pavlov. The asymptotic distribution of maximum tree size in a random forest. *Theor. Probab. Appl.*, 22(3):509–520, 1978.
- [14] Yu. L. Pavlov. Limit distributions of the height of a random forest. *Teor. Veroyatnost. i Primenen.*, 28(3):449–457, 1983.
- [15] Jim Pitman. Enumerations of trees and forests related to branching processes and random walks. In *Microsurveys in discrete probability (Princeton, NJ, 1997)*, volume 41 of *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.*, pages 163–180. Amer. Math. Soc., Providence, RI, 1998.
- [16] Douglas Rizzolo. Scaling limits of Markov branching trees and Galton-Watson trees conditioned on the number of vertices with out-degree in a given set. *arXiv:1105.2528v1*, 2011.
- [17] E. Schröder. Vier combinatorische probleme. *Z. Math. Physik*, 15:361–376, 1870.
- [18] Richard P. Stanley. *Enumerative combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999.

DEPARTMENT OF STATISTICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720
E-mail address: `pitman@stat.berkeley.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720
E-mail address: `drizzolo@math.berkeley.edu`